



# DENTALPRAXIS BERLIN

---

## INFORMED CONSENT FOR SURGICAL TOOTH REMOVAL

### Patient data:

Name: \_\_\_\_\_

Vorname: \_\_\_\_\_

Date of birth: \_\_\_\_\_

I agree that the following tooth/teeth will be removed from me.

\_\_\_\_\_

### Reason for tooth removal:

- Lack of space, proper eruption of the tooth is not guaranteed, pericorony inflammation may occur.
- there is inflammation of the root tip
- the success of orthodontic treatment may be jeopardised
- there is a deep caries that can no longer be treated
- A cyst has developed on the tooth
- the tooth is the suspected cause of an inflammation or cyst of the maxillary sinus
- the dental care of neighbouring teeth is prevented

### The procedure takes place in:

- local anaesthetic

**Additional service:** Socket preservation incl. material (collagen cone): 25, 00 Euro

necessary

Not necessary

Surgical tooth removal involves exposing the tooth through a mucosal incision and removing the bone with a drill. Sometimes it is necessary to grind the tooth before it can be removed with a lever or forceps. The wound is sutured or sometimes filled with a gauze strip for drainage.

---

### Dentalpraxis Berlin

Ferda Betül Igac **IBAN:** DE19 3006 0601 0029 8955 07  
Fuggerstr. 35 **BIC:** DAAEDEDXXX  
10777 Berlin

### Bank details: Deutsche Apotheker- und Ärztebank Berlin



# DENTALPRAXIS BERLIN

---

Alternative to tooth removal:

---

---

- There are no alternatives

Are complications to be expected?

In individual cases, complications may occur depending on your health condition, age, lifestyle (smoking) and other factors. Complications can occur during and/or after the operation.

There is a risk of nerve damage, or nerve injury. The consequence can be a permanent restriction of the quality of life that impairs the way of life and can be associated with deficits and discomfort.

General complications: Nervousness, restlessness, drop in blood pressure also due to intolerance to the local anaesthetic. Allergies must be treated immediately as they can be life-threatening.

Bleeding: In case of postoperative bleeding, biting on a clean handkerchief or swab is usually sufficient. If not, further measures will be taken in the practice.

Scarring: If a cut has to be made, scarring will occur.

Inflammation: Treatment with antibiotics, irrigation or a follow-up operation may be necessary.

Special complications: Break-off of fine root tips, depending on the findings they are removed or left in place.

Damage to adjacent tissues by surgical instruments.

Upper jaw: maxillary sinusitis. When broken root tips are opened or penetrated, maxillary sinusitis may occur; treatment with antibiotics may be necessary.

---

**Dentalpraxis Berlin**

Ferda Betül Igac **IBAN:** DE19 3006 0601 0029 8955 07  
Fuggerstr. 35 **BIC:** DAAEDEDXXX  
10777 Berlin

**Bank details:** Deutsche Apotheker- und Ärztebank Berlin



# DENTALPRAXIS BERLIN

---

**Lower jaw:** Sensory disturbances in the area of the lower jaw and lower lip. If the roots of the removed tooth are very close to the mandibular nerve, mechanical damage (temporary or permanent) to the nerve may occur during tooth removal. Also rare are disturbances of sensation or taste in the area of the tongue. These can only occur with the removal of lower wisdom teeth.

**Lower jaw fracture:** If the position of a lower wisdom tooth is unfavourable, the lower jaw may fracture even weeks after the tooth has been removed.

**Bone inflammation:** this can also occur later if the wound does not heal, sometimes with numbness and pain sensation in the affected area. A follow-up operation and/or treatment with antibiotics may be necessary.

**What medication are you currently taking (anticoagulants)?**

---

Do you have any questions or comments?

---

I confirm that I have understood the above information.

All my questions have been answered. I am aware that I can withdraw my consent.

---

**Dentalpraxis Berlin**

Ferda Betül Igac **IBAN:** DE19 3006 0601 0029 8955 07  
Fuggerstr. 35 **BIC:** DAAEDEDXXX  
10777 Berlin

**Bank details:** Deutsche Apotheker- und Ärztebank Berlin